526 Rec'd PCT/PTO Attorney's Docket Number	30JUNZUUU
045636-5037	_

## TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US)

CONCERNING A FILING UNDER 35 U.S.C. 371 International Application. No. International Filing Date Priority Date Claimed

PCT/FR98/02899 Title of Invention

29 December 1998 30 December 1997

#### PEPTIDE EPITOPES RECOGNIZED BY ANTI-FILAGGRIN AUTO-ANTIBODIES IN SERUM FROM RHEUMATOID ARTHRITIS PATIENTS

ng items and other

	App	licants Fo	r DO/EO	<i>r</i> us		
	DA	LBON, Co	lette JOI	GIRBAL-NEUHAUSER, Christian VINCENT, Michel SIMON, Mireille SEBBAG, F JVET-REYNAUD and Michel ARNAUD submit to the United States Designated/Elected Office (DO/EO/US) the following iter		
_		mu i		T		
1.	[X]			T submission of items concerning a filing under 35 U.S.C. 371.		
2.	[ ]			OND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371.		
3.	[]	than	This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(l).			
à				and for International Preliminary Examination was made by the 19th month from the		
4.	[X]			I priority date.		
5.	₹₫ [X]			International Application as filed (35 U.S.C. 371(c)(2))		
٥.		a.	[]	is transmitted herewith (required only if not transmitted by the		
	14			International Bureau).		
	14	b.	[X]	has been transmitted by the International Bureau.		
	ļā ļā	c.		is not required, as the application was filed in the United States Receiving Office (RO/US).		
6.	* [X]	A tra	nslation o	f the International Application into English (35 U.S.C. 371(c)(2)).		
7.	[n [X]			o the claims of the International Application under PCT Article 19		
		(35 L	J.S.C. 37			
	13	a.	[]	are transmitted herewith (required only if not transmitted by the International Bureau).		
	(3	b.	[]	have been transmitted by the International Bureau.		
	(3	c.	[]	have not been made; however, the time limit for making such amendments has NOT expired.		
		d.	[X]	have not been made and will not be made.		
8.	[ ]			f the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).		
9.	[ ]			elaration of the inventors (35 U.S.C. 371(c)(4)).		
10.	[]			f the annexes to the International Preliminary Examination Report ticle 36 (35 U.S.C. 371(c)(5)).		
Iter	ns 11. to 1	6. below	concern o	other document(s) or information included:		
11.	[X]			n Disclosure Statement under 37 CFR 1.97 and 1.98.		
12.	[]			document for recording. A separate cover sheet in compliance with		
				and 3.31 is included.		
13.	[X]			minary amendment.		
	[ ]			r SUBSEQUENT preliminary amendment.		
14.	[ ]			ecification.		
15.	[]			tement Claiming Small Entity Status		
16.	[X]			information:		
		a.	[X]	Copy of Form PCT/IB/304		
		b. с.	[X]	Copy of Form PCT/IB/308 WIPO Publication WO 99/35167		

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The | assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property | \$0.00 |

TOTAL FEES ENCLOSED = \$840.00 |
Amount to be refunded | \$

a. [X] A check in the amount of \$840.00 to cover the Basic fee is enclosed. b. [] Please charge my Deposit Account No. 50-0310 in the amount of \$0.00

to cover the above fees. A duplicate copy of this sheet is enclosed.

c. [X] Except for issue fees payable under 37 C.F.R. § 1.18, the Commissioner is hereby authorized by this paper to charge any additional fees during the entire pendency of this application including fees due under 37 CFR § 1.16 and

§1.17 which may be required, or credit any overpayment to Deposit Account No. 50-0310.

SEND ALL CORRESPONDENCE TO: Morgan, Lewis & Bockius LLP 1800 M Street, N.W. Washington, D.C. 20036 (202) 467-7000 Elizabeth C. Weimar Rep. No. 44.478

charged

1.8

a weigh

Submitted: June 30, 2000

# 09/582711 532 Rec'd PCT/PTC 30 JUN 2000

PATENT ATTORNEY DOCKET NO. 045636-5037

#### IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re A	application of	f: SERRE et al.	)	
U.S. A	pplication N	fo.: To Be Assigned	)	Group Art Unit: Unassigned
Date o	f National		)	
Stage	Entry	: June 30, 2000	)	Examiner: Unassigned
Based	on PCT/FR	98/02899	)	
Filed		: December 29, 1998	)	
			)	
For:	PEPTIDE E	EPITOPES RECOGNIZED BY	( )	
	ANTI- FIL.	AGGRIN AUTOANTIBODIE	S )	
	IN SERUM	FROM RHEUMATOID	)	
	ARTHRITI	S PATIENTS	)	
	ant Commiss ngton, D.C.	sioner for Patents 20231		
Sir:				

#### PRELIMINARY AMENDMENT

Prior to the examination of the above-identified application on the merits, please amend the application, without prejudice, as follows:

ABSTRACT: Please add the abstract provided on a separate sheet.

### IN THE CLAIMS:

Please cancel claim 4.

#### Please amend claims 1-3 and 5-7 as follows:

Claim 1, line 1, change "Peptide" to -- A peptide--;

Claim 1, line 3, change "characterized in that" to -- wherein --;

Claim 1, line 4, change "centred" to -- centered --;

Claim 2, line 1, change "Peptide" to -- The peptide--; and, change "characterized in" to -- wherein --;

Claim 2, line 2, delete "that it comprises";

after "motif" add -- comprises --; and,

after "Ser-Cit-His" add a comma -- , --;

Claim 3, line 1, change "Artificial" to -- An artificial --;

Claim 3, line 3, delete "characterized in that it";

Claim 3, line 4, change "comprises" to -- comprising --; and delete "or consists of";

2

1-WA/1437762.1

```
Claim 3, line 5, change "either of Claims 1 and 2" to -- Claim 1 --;
```

Claim 5, line 1, change "Antigenic" to -- An antigenic --;

Claim 5, line 3, delete "characterized in that it";

Claim 5. line 4, change "contains" to -- comprising --; and, delete "any one of";

Claim 5, line 5, change "Claims" to -- Claim --; and,

delete the remainder of the line, "to 3 which is optionally labelled with

and/or";

Claim 5, line 6, delete the entire line except the period, "conjugated to a carrier molecule";

Claim 6, line 1, change "Method" to -- A method --;

Claim 6, line 2, delete "this";

Claim 6, line 3, delete "method being characterized in that it"; and, change "comprises" to - comprising --;

```
Claim 6, line 5, change "any one of Claims 1 to 3" to -- Claim 1 --;
```

Claim 6, line 7, change "the" to -- any --;

Claim 6, line 8, delete "possibly"; and,

after "present" add -- in the biological sample --;

Claim 6, line 9, change "the" to -- any --;

Claim 6, line 10, delete "possibly";

Claim 7, line 1, change "Kit" to -- A kit --;

Claim 7, line 2, delete "characterized in";

Claim 7, line 3, change "that it comprises" to -- comprising --; and, delete "any";

Claim 7, line 4, change "one of Claims 1 to 3" to -- Claim 1 --;

Claim 7, lines 7 and 8, delete in their entirety except the period, "and/or means for detecting said antigen/antibody complex".

# Please add the following claims 8-12.

- 8. The artificial antigen of Claim 3 wherein the antigen consists of at least one peptide comprising a tripeptide motif centered on a citrulline residue, which is specifically present on at least one of the citrullinated peptides derived from the sequences SEQ ID NO: 3, SEQ ID NO: 5 or SEO ID NO: 6.
  - 9. The antigenic composition of Claim 5 wherein the at least one antigen is labeled.
- 10. The antigenic composition of Claim 5 wherein the at least one antigen is conjugated to a carrier molecule.
  - 11. The kit of Claim 7 further comprising means for detecting the antigen/antibody complex.
- 12. A kit for detecting rheumatoid arthritis-specific autoantibodies in a biological sample, comprising at least one antigen according to Claim 1, as well as means for detecting the antigen/antibody complex.

5

1-WA/1437762.1

#### REMARKS

The changes to the claims requested above have been made so as to eliminate multiple claim dependencies and to present claim language more conventional for practice in the United States. These changes do not introduce new matter, nor do they alter the subject matter presented and examined in the corresponding International Application.

Respectfully submitted,

MORGAN, LEWIS & BOCKIUS LLP

By: Elizabeth C. Volomor Elizabeth C. Weimar

Reg. No. 44,478

Dated: June 30, 2000 MORGAN, LEWIS & BOCKIUS LLP Intellectual Property Support Unit 1800 M Street, NW Washington, DC 20036-5869

(202) 467-7000

# ABSTRACT OF THE DISCLOSURE

The invention concerns peptides comprising epitopes recognized by antifilaggrin in serum from rheumatoid arthritis patients. Said epitopes comprising a tripeptide motif centered on a citrulline residue. The invention also concerns artificial antigens comprising said epitopes, and their use for diagnosing rheumatoid arthritis.

# 532 Rec'd PCT/PTC 3 0 JUN 2000 PCT/FR98/02899

PEPTIDE EPITOPES RECOGNIZED BY ANTI-FILAGGRIN AUTO-ANTIBODIES IN SERUM FROM RHEUMATOID ARTHRITIS PATIENTS

The present invention relates to novel preparations of antigens which are specifically recognized by rheumatoid arthritis-specific autoantibodies.

Rheumatoid arthritis (hereafter abbreviated to "RA") is the most frequent of the chronic inflammatory rheumatisms. It is an autoimmune disease, and the serum from affected patients contains autoantibodies, certain of which are specific, and may constitute a marker for this disease, allowing its diagnosis even at early stages. Research has thus been carried out with a view to identifying antigens recognized by these antibodies, in order to obtain purified preparations thereof which can be used in conventional techniques of immunological diagnosis.

Autoantibodies which are specifically present in RA patients, and which react with a rat oesophagial epithelial antigen, were described for the first time by B.J.J. Young et al. in Br. Med. J. 2:97-99, (1979). These autoantibodies were, at the time, named "antikeratin antibodies".

During previous work, the inventors' team obtained, from human and murine malpighian epithelia, 25 preparations of antigens related to filaggrin and to profilaggrin which are recognized specifically by antibodies present in serum from rheumatoid arthritis patients, and showed that the "anti-keratin antibodies" were in fact anti-filaggrin autoantibodies (hereafter named "AFA"). Application EP 0 511 116 describes these antigenic preparations and their use for diagnosing rheumatoid arthritis.

Filaggrins are a family of proteins which have been identified in various species, inter alia in humans, rats, mice and guinea pigs, in keratinizing malpighian epithelia [for a review on filaggrins, cf. Dale et al. [The Keratinocyte Handbook, Cambridge

10

15

20

25

30

35

University Press, pp. 323-350, (1994)]]. They are derived from the dephosphorylation and proteolysis of a precursor, profilaggrin, which consists essentially of repeated filaggrin domains separated by interdomain peptide segments.

The gene encoding profilaggrin is composed of repeated subunits, each of which encodes a filaggrin molecule, separated by portions encoding the interdomain peptide segments. All the repeat units encoding each of the human filaggrins have the same length (972 base pairs in humans); however, in humans, considerable sequence variation (10-15%) is observed from one subunit to the other. While most are conservative, some of these variations induce amino acid changes and, in certain cases, changes in the electrical charge of the protein. Thus, human filaggrins form, independently of post-transcriptional modifications, a heterogeneous population of molecules of similar size, but of different sequences and charges (pHi equal to 8.3 ± 1.1) [Gan et al., Biochem. 29, pp. 9432-9440 (1990)].

Profilaggrin is a protein of high molecular weight (approximately 400,000 in humans) which is soluble in the presence of high concentrations of salts or of urea. It possesses a high content of basic amino acids (arginine and histidine) and of glycine, serine and glutamic acid. It is low in nonpolar amino acids, and contains neither methionine, nor cysteine, nor tryptophan. It is highly phosphorylated on serine residues, which gives it an isoelectric point close to neutrality.

Profilaggrin is cleaved into filaggrin units during a complex maturation process which involves dephosphorylation, followed by cleavage by proteases at the interdomain segments. This cleavage generates, first of all, fragments of intermediate size, and then the functional filaggrin molecules.

Filaggrins derived from the dephosphorylation and cleavage of profilaggrin are basic proteins, the amino acid content of which is similar to that of

10

15

20

25

30

35

profilaggrins. They participate in the organization of keratin filaments, and undergo a progressive maturation during which the arginine residues, which are basic, are converted into citrulline residues, which are neutral, under the action of peptidylarginine deiminase [Harding C.R. and Scott I.R., J. Mol. Biol. 170, pp. 651-673 (1983)]. This leads to a reduction in their affinity for keratins, from which they detach; they are then totally degraded under the action of various proceases.

Filaggrin and profilaggrin properties have been particularly well studied in rats, in mice and in humans. The size of profilaggrin varies, according to the species, from 300 to 400 kD, and that of filaggrins from 27 to 64 kD.

The polymorphism observed in humans between the filaggrin units within t.he profilaggrin gene does not appear in rats and mice. Filaggrins also exhibit great inter- and intraspecific variability in their sequence. This variability does not however affect their functional properties, or their overall amino acid composition, or their biochemical properties. Similarly, the localizations of profilaggrin and of filaggrins are identical in the various mammals studied.

In continuing their work, the inventors noticed that profilaggrin present in keratohyalin granules of human epidermis was not recognized, unlike filaggrins, by AFAs [Simon et al. Clin. Exp. Immunol. 100, 90-98 (1995)]. They then tested the reactivity of AFAs with recombinant filaggrin, and observed that this was not recognized either. Conversely, it had previously been observed that the forms of human epidermal filaggrin principally recognized by AFAs were the acido-neutral forms described by Simon et al. [J. Clin. Invest., 92, 1387, (1993)], and in Application EP 0 511 116. The fact that these acido-neutral forms correspond to a late maturation stage of filaggrin made it possible to presume that all or part of the post-translational

1.5

20

25

30

modifications which intervene up to this stage are involved in the formation of the epitopes recognized by AFAs.

To verify this hypothesis, the inventors attempted to reproduce these post-translational modifications in vitro, using recombinant filaggrin, in order to determine which are capable of influencing filaggrin antigenicity.

They thus in fact observed that citrullination of filaggrin was enough to generate epitopes recognized by AFAs. Specifically, they observed, by carrying out in vitro deimination of recombinant filaggrin, that replacing at least one portion of the arginines with citrullines allows the production of an antigen which is recognized specifically by AFAs present in serum from RA patients. They also located regions which, after citrullination, were highly immunoreactive with respect to anti-filaggrin autoantibodies. They are in particular the region corresponding to the C-terminal portion (amino acids 144 to 324), and in particular to amino acids 144 to 314, as well as the region corresponding to amino acids 76 to 144 and the region corresponding to amino acids 71 to 119, of a human filaggrin unit. This work resulted in the production of artificial antigens, which are recognized specifically by AFAs present in serum from RA patients, and which consist of recombinant or synthetic polypeptides derived from the sequence of filaggrin or from portions of it, by substituting at least one arginine residue with a citrulline residue. These antigens, as well as their use, form the subject of Application FR 96/10651, filed on 30 August 1996 in the name of Biomérieux.

In continuing their work, the inventors managed to select, using the sequence of one filaggrin unit, septides in which substituting at least one arginine residue with a citrulline residue gives rise to epitopes which are recognized specifically by AFAs present in serum from RA patients.

The sequences of these peptides are identified

15

20

25

35

in the attached sequence listing under the numbers SEQ ID NO: 3, SEQ ID NO: 5 and SEQ ID NO: 6.

"Filaggrin unit" is intended to mean a polypeptide, the sequence of which is that of the translation product of any one of the subunits encoding a filaggrin domain of the human, or any other species, profilaggrin gene, or is a consensus sequence, this theoretical sequence being obtained from filaggrin domain sequences.

The inventors have now identified epitopes recognized by anti-filaggrin autoantibodies: these epitopes comprise a tripeptide motif centred on a citrulline residue, which is specifically present on the citrullinated peptides derived from the sequences SEQ ID NO: 3, SEQ ID NO: 5 or SEQ ID NO: 6, and which is absent from the sequence SEO ID NO: 4.

It is in particular the tripeptide motif Ser-Cit-His in which Cit represents a citrulline residue.

A subject of the present invention is a peptide which constitutes an epitope recognized by antifilaggrin autoantibodies present in serum from rheumatoid arthritis patients, characterized in that said epitope comprises a tripeptide motif centred on a citrulline residue, which is specifically present on at least one of the citrullinated peptides derived from the sequences SEQ ID NO: 3, SEQ ID NO: 5 or SEQ ID NO: 5.

According to a preferred embodiment of the present invention, said peptide comprises at least one pentapeptide motif centred on a citrulline residue, which is present on at least one of the citrullinated peptides derived from the sequences SEQ ID NO: 3, SEQ ID NO: 5 or SEQ ID NO: 6.

Advantageously, said peptide comprises the tripeptide motif Ser-Cit-His in which Cit represents a citrulline residue.

By way of example, mention will be made of peptides derived, by citrullination, from peptides

15

20

25

which comprise the pentapeptide motif X1-Ser-Arg-His-X2 in which X1 = Ser or Gly, and X2 = Ser or Pro, and among them, peptides which comprise the hexapeptide motif X0-X1-Ser-Arg-His-X2 or the heptapeptide motif X0-X1-Ser-Arg-His-X2-X3, in which X1 and X2 are as defined above, X0 = Asp and X3 = Gly or Arg.

The peptides in accordance with the invention allow the preparation of artificial antigens which are recognized specifically by anti-filaggrin autoantibodies present in serum from rheumatoid arthritis patients. These artificial antigens also form part of the subject of the present invention.

Artificial antigens in accordance with the invention comprise at least one peptide epitope centred on a citrulline residue as defined above. They consist for example of peptides of at least 5 amino acids, preferably at least 10 amino acids, and advantageously at least 14 amino acids. They can be peptides consisting of at least one fragment of at least one of the citrullinated peptides derived from the sequences SEQ ID NO: 3, SEQ ID NO: 5 or SEQ ID NO: 6, or containing at least one such fragment. These peptides can comprise several citrullinated epitopes which are specifically recognized by AFAs, and which are of identical or different sequences.

The term "peptide" as used in the present application means in particular protein or protein oligopeptide, which is extracted, fragment, or separated or substantially isolated or synthesized, in particular those obtained by chemical synthesis or by 30 expression in a recombinant organism; any peptide in the sequence of which one or more amino acids of the L series are replaced with an amino acid of the D series, and vice versa; any peptide in which one at least of the CO-NH bonds, and advantageously all the 35 CO-NH bonds, of the peptide chain is (are) replaced with one or more NH-CO bonds; any peptide in which one at least of the CO-NH bonds, and advantageously all the CO-NH bonds, is or are replaced with one or more NH-CO

15

20

25

30

35

bonds, the chirality of each aminoacyl residue, whether or not it is involved in one or more abovementioned CO-NH bonds, being either conserved or inverted with respect to the aminoacyl residues constituting a reference peptide, these compounds also being referred to as immunoretroids, a mimotope, etc.

Antigens in accordance with the invention can for example be obtained by reacting PAD (peptidylarginine deiminase) with natural, recombinant or synthetic proteins or peptides comprising arginine residues and in particular comprising at least one arginine residue constituting the centre of a tripeptide motif identical to those present in sequences SEQ ID NO: 3, SEQ ID NO: 5 and SEQ ID NO: 6; they can also be obtained by peptide synthesis, by directly incorporating one or more citrulline residues, and preferably one or more epitopes comprising a defined above, into citrulline residue as synthesized peptide.

A subject of the present invention is also the use of the antigens in accordance with the invention as defined above for diagnosing RA in vitro.

The present invention encompasses in particular antigenic compositions for diagnosing the presence of RA-specific autoantibodies in a biological sample, these compositions being characterized in that they contain at least one antigen in accordance with the invention which is optionally labelled with and/or conjugated to a carrier molecule.

A subject of the present invention is also a method for detecting RA-specific class G autoantibodies in a biological sample, this method being characterized in that it comprises:

- bringing said biological sample into contact with at least one antigen in accordance with the invention as defined above under conditions which allow the formation of an antigen/antibody complex with the RA-specific autoantibodies possibly present;
  - detecting, by any suitable means, the antigen/anti-

body complex possibly formed.

This detection method can be implemented using a kit comprising at least one antigen according to the invention, as well as suitable buffers and reagents for constituting a reaction medium which allows the formation of an antigen/antibody complex, and/or means for detecting said antigen/antibody complex.

Said kit can also comprise, where appropriate, reference samples such as one or more negative serum (sera) and one or more positive serum (sera).

The present invention will be better understood with the aid of the further description which will follow, which refers to examples of preparation and of use of antigens in accordance with the invention.

# EXAMPLE 1: IN VITRO DEIMINATION OF RECOMBINANT FILAGGRIN WITH PEPTIDYLARGININE DEIMINASE (P.A.D.)

A DNA fragment encoding a filaggrin unit is amplified from human genomic DNA (RAJI cells: ATCC CCL86) by PCR using the 2 primers below:

- 5' primer:
- 5' TTCCTATACCAGGTGAGCACTCAT 3'
- 3' primer:
- 5' AGACCCTGAACGTCCAGACCGTCCC 3'

The amplification product is cloned into the SmaI site of the vector pUC19. The recombinant clones are selected by verifying the presence of a 972-bp insert obtained after digestion with SacI and XbaI. This insert is then subcloned into pUC19. The insert resulting from this subcloning is then transferred into the vector pGEX (sold by the company Pharmacia), between the EcoRI and HindIII sites. In E. coli, the expression vector thus obtained expresses filaggrin in fusion with glutathione-S-transferase (GST) under the control of the Tac prokaryotic promoter. Recombinant protein synthesis is induced by adding isopropyl- $\beta$ -D-galactoside (IPTG) to the culture.

15

20

25

30

35

1.0

The recombinant filaggrin thus obtained will be named hereafter: "fil-gst".

After electrophoresis, the existence of 9 fragments is observed, which result from a post-translational proteolysis of whole filaggrin.

The mixture of the 9 fragments is subjected to an *in vitro* deimination with peptidylarginine deiminase.

A preparation of rabbit muscle peptidylarginine 10 deiminase (681 U/ml), sold by Takara Biomed Europe, is used according to the protocol recommended by the manufacturer.

The procedure conditions are as follows:

- reaction medium: 0.1 M Tris-HCl, 10 mM CaCl2,
- 15 5 mM DTT, pH 7.4;
  - enzyme/substrate ratio: 140 mU/ $\mu$ mol of filaggrin containing 10% of arginine, i.e. 4 mU/ $\mu$ mol of arginine;
    - incubation: between 0 and 60 min at 50°C;
  - termination of the reaction: heating for 3 min in Laemmli buffer.

The 8 reactions below are carried out in parallel.

- (1) BSA (bovine serum albumin) incubated in reaction medium (1 h, 50°C) without P.A.D.
  - (2) BSA incubated in reaction medium (1 h, 50°C) with 60 mU of P.A.D.
  - (3) fil-gst incubated in reaction medium (1 h, 50°C) without P.A.D.
- 30 (4) fil-gst incubated in reaction medium (5 minutes at 50°C) with 60 mU of P.A.D.
  - (5) fil-gst incubated in reaction medium (15 minutes at 50°C) with 60 mU of P.A.D.
- (6) fil-gst incubated in reaction medium 35 (30 minutes at 50°C) with 60 mU of P.A.D.
  - (7) fil-gst incubated in reaction medium (1 h at 50°C) with 60 mU of P.A.D.
  - (8) fil-gst incubated in reaction medium (1 h at 50°C) with 60 mU of P.A.D. and in the presence of

15

20

25

35

10 mM of N-ethylmaleimide (inhibitor of P.A.D.).

1  $\mu$ l of each sample is loaded on an electrophoresis gel (12.5% SDS-PHAST® gel, Pharmacia), and electrophoresis is carried out using the PHAST-SYSTEM® apparatus (Pharmacia) under the conditions recommended by the manufacturer. After transfer onto nitrocellulose, detection is carried out either with a pool of 5 sera from RA patients, diluted to 1/2000, or with the anti-filaggrin monoclonal antibody AHF2 [Simon et al. J. Invest. Dermatol. 105, 432, (1995)] at the concentration of 0.2  $\mu$ G/ml.

The antigen/antibody complex is detected by the ECL technique using a peroxidase-coupled secondary antibody.

The results show that, in the absence of citrullination reaction, the fil-gst is not recognized by sera from RA patients, whereas right from 5 minutes of citrullination, it is detected by these sera. An increase in the reactivity with the pool of sera is observed when P.A.D. is reacted for 60 minutes at 50°C.

In addition, these results make it possible to presume that one or more epitopes of high affinity exist in the COOH-terminal moiety of filaggrin (positions 144 to 324), this or these epitope(s) being repeated between positions 76 and 144.

# EXAMPLE 2: CITRULLINATION OF PEPTIDES S-47-S AND S-35-R WITH P.A.D., AND CITRULLINATED PEPTIDE REACTIVITY ASSAY

The 49-amino acid peptide S-47-S of sequence 30 (1-letter code):

 ${\tt NH_2-STGHSGSQHSHTTTQGRSDASRGSSGSRSTSRETRDQEQSGDGSRHSGS-COOH}$ 

corresponding to amino acids 71 to 119 of the sequence of a human filaggrin unit and comprising 6 arginine residues, and

the 37-amino acid peptide S-35-R of sequence (1-letter code):

NH2-SQDRDSQAQSEDSERRSASASRNHRGSAQEQSRDGSR-COOH

corresponding to amino acids 155 to 191 of the sequence of a human filaggrin unit and comprising 7

10

15

20

25

arginine residues were prepared by peptide synthesis. The peptides S-47-S and S-35-R are represented in the attached sequence listing under the respective numbers SEQ ID No: 3 and SEQ ID No: 4.

These 2 peptides, as well as fil-gst, were citrullinated by reacting P.A.D. for 30 minutes at  $50^{\circ}$ C in the same reaction medium as that indicated in Example 1. The conditions specific for each peptide and for fil-gst are as follows:

- peptide S-47-S: 4 mU/µmol arginine
  - peptide S-35-R: 2.7 mU/ $\mu$ mol arginine
  - fil-gst: as indicated in Example 1.

The reactivity of each peptide and that of fil-gst, before and after reacting the enzyme, with respect to the monoclonal antibody AHF4 and to the serum from an RA patient are compared by "dot-blot".

The procedure conditions are as follows:

- 0.5  $\mu$ g per deposit of each antigen (peptides, fil-gst, acido-neutral variants of filaggrin (AVF)).
- Treatment of the nitrocellulose for  $45 \text{ minutes at } 80 \, ^{\circ}\text{C}$ , before immunodetection.
- RA serum used at the dilution of 1/2000; monoclonal antibody AHF4 used at a concentration of 0.2  $\mu g/ml$ .

The results show that:

- AHF4 recognizes citrullinated or noncitrullinated peptide S-47-S and fil-gst, but does not recognize citrullinated or noncitrullinated S-35-R.
- S-47-S is recognized, after citrullination, 30 by the serum of the RA patient, whereas neither citrullinated nor noncitrullinated S-35-R is recognized. The same serum recognizes moreover citrullinated fil-gst and AVFs, but does not recognize noncitrullinated fil-gst.

15

# EXAMPLE 3: SYNTHESIS OF CITRULLINATED AND NON-CITRULLINATED PEPTIDES E-12-H AND E-12-D, AND PEPTIDE REACTIVITY ASSAY

The peptides E-12-H and E-12-D were determined by reference to the nucleotide sequences of the human profilaggrin gene described by Gan S.Q. et al. [Biochemistry, 29: 9432-9440, (1990)].

The 14-amino acid peptide E-12-H of sequence (1-letter code):

NH2-EQSADSSRHSGSGH-COOH

comprises 1 arginine residue, and

the 14-amino acid peptide E-12-D of sequence (1-letter code):

NH2-ESSRDGSRHPRSHD-COOH

comprises 3 arginine residues.

The peptides E-12-H and E-12-D are represented in the attached sequence listing under the respective numbers SEQ ID No: 5 and SEQ ID No: 6.

 $\qquad \qquad \text{These peptides were prepared by solid phase} \\ 20 \quad \text{peptide synthesis.}$ 

The citrullinated peptides E-12-H and E-12-D were synthesized directly by incorporating a citrulline as a replacement for an arginine.

For the peptide E-12-D, only the arginine 25 residue corresponding to the 8<sup>th</sup> amino acid of the sequence was replaced with a citrulline during the peptide synthesis.

The reactivity of each citrullinated and non-citrullinated peptide was assayed with respect to a normal serum, to two sera from RA patients, to antifilaggrin antibodies (AFAs) purified from a pool of 45 sera from RA patients and to anti-filaggrin antibodies purified from 12 sera from RA patients, respectively.

35

30

#### EXPERIMENTAL PROTOCOL:

The wells of Nunc Maxisorp microtitration plates were covered using the noncitrullinated and citrullinated peptides E-12-D and E-12-H, respectively,

15

20

diluted to a concentration of 5  $\mu$ g/ml in a PBS buffer (pH: 7.4) and incubated overnight at 4°C (final volume: 100 µl/well). The wells were saturated for 30 minutes at 37°C with 200 µl/well of PBS-Tween 20 (0.05%), 2.5% gelatin. The negative control serum (normal serum) was diluted to 1/120. The anti-filaggrin antibodies were diluted in PBS-Tween 20 (0.05%)-gelatin (0.5%) (PBS TG) such that the final concentrations of anti-filaggrin autoantibodies are those indicated in the attached Table I. The negative control serum, the RA sera and the anti-filaggrin antibodies were added (final volume: 100  $\mu$ l/well) and subjected to incubation for 1 hour at 37°C and overnight at 4°C. Peroxidase-labelled goat anti-human immunoglobulin gamma heavy chain antibodies (sold by the company Southern Biotechnologies) were added to each well (dilution in PBSTG: 1/2000, final volume: 100 µl/well) and subjected to incubation for 1 hour at 37°C. The revelation was carried out by ortho-phenylenediamine (2 mg/ml, for adding 10 minutes).

The results presented in the attached Table I are given as a ratio of OD at 492 nm: citrullinated peptide signal/noncitrullinated peptide signal.

These results show that, in the majority of cases, the citrullinated peptide/noncitrullinated peptide OD ratio is higher than 1, and thus illustrate the good sensitivity of the citrullinated peptides, compared to the noncitrullinated peptides, as regards their reactivity with respect to anti-filaggrin autoantibodies.

# TABLE I

ſ		<u>*</u>	92	84
		10	1 1.	2 5.
1		10*	1.1	1.2
		10*	1.57	3.66
	sera	10*	1.13	1.14
	.2 RA	10*	5.19	3.18
	[rom]	10*	1.87	4.14
	AFAs purified from 12 RA sera	10*	3.50	3.57
	puri	10*	1.12	0.76 1
	AFAS	10*	1.19	1.18
		10*	2.48	1.06
		10.	1.38	2.05
		10.	1.99	7.16
	51	20*	1.48	1.82
	AFA pool	10*	1.63	2.45
	AE	2*	1.11	2.45
	~	*02	5.57	8.38
	erum	10*	1.1.8	1,51
	RA serum 2	5.	. 42	1.32 1.20 10.44 11,51 8.38 2.45 2.42 1.02 7.16 2.05 1.06 1.18 0.76 13.57 4.14 3.18 1.14 3.66 1.22 5.84
		*	85 2	20 1
	serm	20	2 1.	2 1.
	2	10.	1.4	1.3
	Control	serum 10* 20* 5* 10* 20* 5* 10* 20* 10* 10* 10* 10* 10* 10* 10* 10* 10* 1	1.076	1
	Peptide Control RA serum 1		E-12-D 1.076 1.42 1.68 2.42 3.77 5.57 1.77 1.63 1.48 1.99 1.38 2.48 1.19 1.12 3.50 1.87 5.19 1.13 1.57 1.11 1.65	Е-12-Н

\*: AFA concentration in µg/ml.

1.0

15

25

30

35

#### CLATMS

- 1) Peptide comprising an epitope recognized by anti-filaggrin autoantibodies present in serum from rheumatoid arthritis patients, characterized in that said epitope comprises a tripeptide motif centred on a citrulline residue, which is specifically present on at least one of the citrullinated peptides derived from the sequences SEQ ID NO: 3, SEQ ID NO: 5 or SEQ ID NO: 6.
  - 2) Peptide according to Claim 1, characterized in that it comprises the tripeptide motif Ser-Cit-His in which Cit represents a citrulline residue.
- 3) Artificial antigen recognized specifically by anti-filaggrin autoantibodies present in serum from rheumatoid arthritis patients, characterized in that it comprises or consists of at least one peptide according to either of Claims 1 and 2.
- 4) Use of an antigen according to any one of 20 Claims 1 to 3 for diagnosing rheumatoid arthritis in vitro.
  - 5) Antigenic composition for diagnosing the presence of rheumatoid arthritis-specific autoantibodies in a biological sample, characterized in that it contains at least one antigen according to any one of Claims 1 to 3 which is optionally labelled with and/or conjugated to a carrier molecule.
  - 6) Method for detecting rheumatoid arthritisspecific autoantibodies in a biological sample, this method being characterized in that it comprises:
  - bringing said biological sample into contact with at least one antigen according to any one of Claims î to 3 under conditions which allow the formation of an antigen/antibody complex with the rheumatoid arthritis-specific autoantibodies possibly present;
  - detecting, by any suitable means, the antigen/antibody complex possibly formed.
  - 7) Kit for detecting rheumatoid arthritis-specific autoantibodies in a biological sample, characterized in

that it comprises at least one antigen according to any one of Claims 1 to 3, as well as suitable buffers and reagents for constituting a reaction medium which allows the formation of an antigen/antibody complex, and/or means for detecting said antigen/antibody complex.

# COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY

U.S. DEPARTMENT OF COMMERCE Patent and Trademark Office

ATTORNEY DOCKET NO.: 045636-5037

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,			
I believe I am the original, first and sole inventor (if only one name is listed below) of an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:			
PEPTIDE EPITOPES RECOGNIZED BY ANTI-FILAGGRIN AUTO-ANTIBODIES IN SERUM FROM RHEUMATOID ARTHRITIS PATIENTS			
the specification of which:			
is attached hereto; or			
was filed as United States application Serial No on and was amended on (if applicable); or			
was filed as PCT international application Number PCT/FR98/02899 on 29 December 1998, and was amended under PCT Article 19 on (if applicable).			
I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.			
I acknowledge the duty to disclose to the U.S. Patent and Trademark Office information which is material to the patentability of claims presented in this application in accordance with Title 37, Code of Federal Regulations, §1.56.			
Thereby claim foreign priority benefits under Title 35, United States Code, §119(a)-(d) or §365(b) of any foreign application(s) for patent or invarious configuration of 335(a) of any PCT international application(s) designating at least one country other than the United States of			

I hereby claim foreign priority benefits under Title 35, United States Code, \$119(a)-(d) or \$365(b) of any foreign application(s) for patent or inventor's certificate or \$365(a) of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a

filing date before that of the application(s) of which priority is claimed:

PRIOR FOREIGN APPLICATION(S):				
COUNTRY (if PCT, indicate PCT)	APPLICATION NUMBER	DATE OF FILING (day, month, year)	PRIORITY CLAIMED	
France	FR 97 16,673	30 December 1997	[X] Yes [ ] No	
			[]Yes []No	

1-84/1437413 1

Page 1 of 6

	nal Applications)		ATTORNEY DOC	KET NO.: 045636-50
I hereby claim the benefits u listed below.	nder Title 35, United States C	ode §119(e) of any Unit	ed States provisional a	pplication(s)
	U.S. PROVISION	AL APPLICATIONS		
U.S. PROVISIONAL APP	LICATION NO.		U.S. FILING DATE	
provided by the first paragra Patent and Trademark Offic application in accordance w date of the prior application	claims of this application is n ph of Title 35. United States e all information known to me tith Title 37, Code of Federal F (s) and the national or PCT int ONS OR PCT INTERNATIO	Code, § 112, I acknowled to be material to the pa- tegulations, § 1.56 which emational filing date of	ige the duty to disclose tentability of claims pr h became available bet this application:	resented in this ween the filing
PRIOR U.S. APPLICATION	NAS OR FOLIMITED WITE	tito rationalities		
U.S. APPLICA	TIONS		STATUS (Check On	e)
	U.S. FILING DATE	PATENTED	PENDING	ABANDONED
U.S. APPLICA		PATENTED		
U.S. APPLICA		PATENTED		
U.S. APPLICA		PATENTED		
U.S. APPLICATION NO.  POWER OF ATTORNES	U.S. FILING DATE  (: As a named inventor, I here c Customer Number provided & Office connected therewith,	by appoint the registers	PENDING  ed practitioners of Mor	ABANDONEL  gan, Lewis &  sact all business

1-44/1437413.1

Page 2 of 6

33 01 45 62 04 86 PAGE.05

Fax émis par: 33 01 45 62 04 86 ORES le 25/09/00 15:54 M4 NORM Pg: 6/9

Combined Declaration For Patent Application and Power of Attorney - (Continued) (includes Reference to PCT International Applications)

ATTORNEY DOCKET NO.: 045636-5037

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

FULL NAME OF SOLE OR FIRST INVENTOR	Guy SERRE	
RESIDENCE & CITIZENSHIP	Toulouse, France	COUNTRY OF CITIZENSHIP FRANCE
POST OFFICE ADDRESS	Appartement 46, Résidence du Lac, 10, avenue Winston Church France	nill, F-31100 Toulouse,
First or sole in	VENTOR'S SIGNATURE GY SKRRE	DATE A grav 25"
FULL NAME OF SECOND INVENTOR	Elisabeth GIRBAL-NEUHAUSER	
RESIDENCE & CITIZENSHIP	Toulouse, France	COUNTRY OF CITIZENSHIP FRANCE
POST OFFICE ADDRESS	22, rue Matelache, F-31000 Toulouse, France	م ا
SECOND INVENTO	R'S SIGNATURE ELisabeth Girs Al. NEUHAUSE	n DATE August 25
Į	isting of Inventors Continued on attached page(s) [X] Yes [	] No

1-44/1437413.1

Page 3 of 6

and the same of th		The state of the s	
	n For Patent Application and Power of Attorney o PCT International Applications)	- (Continued)  ATTORNEY DOCKE	T NO - 045636-5037
		MITORIET BOCK	.1 (40 043030-3037
FULL NAME OF THIRD INVENTOR	Christian VINCENT	1	
RESIDENCE & CITIZENSHIP	Lauzerville, France		COUNTRY OF CITIZEN SHIP FRANCE
POST OFFICE ADDRESS	16, avenue de la Saune, F-31650 Lauzerville	France	
THIRD INVENTOR	S SIGNATURE CHIATOM PINCENT		DATE But 28.2000
FULL NAME OF FOURTH INVENTOR	Michel SIMON		•
RESIDENCE & CITIZENSHIP	Belberaud, France		COUNTRY OF CITIZENSHIP FRANCE
POST OFFICE ADDRESS	2, Lot Cance Merle, F-31450 selberauc	i, France	
FOURTH INVENTO	R'S SIGNATURE Thickel Sinoul		DATE hujuk 29, 2000
Lis Lis	sting of Inventors Continued on attached page(s)	[ ] X Yes	.0

1-40/[4]74[],1

Page 4 of 6

Fax émis par: 33 01 45 62 04 86 ORES le 25/09/00 15:54 A4 NORM Pg: 8/9 Combined Declaration For Patent Application and Power of Attorney - (Continued) (includes Reference to PCT International Applications) ATTORNEY DOCKET NO.: 045636-5037 FULL NAME OF Mireille SEBBAG FIFTH INVENTOR RESIDENCE & Toulouse, France COUNTRY OF CITIZENSHIP CITIZENSHIP FRANCE 3, rue A. Fredeau, F-31500 Toulouse, France Mirithe SEBBAG DATE 8/2012000 Pascal DALBON

POST OFFICE ADDRESS FOURTH INVENTOR'S SIGNATURE FULL NAME OF SIXTH INVENTOR RESIDENCE & Lyon, France COUNTRY OF CITIZENSHIP CITIZENSHIP FRANCE POST OFFICE 6, boulevard Jules Favre, F-69006 Lyon, France ADDRESS Pascal DALBON SIXTH INVENTOR'S SIGNATURE DATE 7/26/2000 ColetteJOLIVET-REYNAUD FULL NAME OF SEVENTH **EXINVENTOR** RESIDENCE & Bron, France COUNTRY OF CITIZENSHIP CITIZENSHIP FRANCE POST OFFICE 16. avenue des ColonnesF-69500 Bron, France ADDRESS SEVENTH INVENTOR'S SIGNATURE DATE 7/28/1000

[-44/[4]74(] [

Colute JOLIVET. REYNAND

Listing of Inventors Continued on attached page(s)

Page 5 of 6

[ | Yes

(X) No

	n For Patent Application and Power of Attorney - (Continue o PCT International Applications)  ATTORNE	ed) Y DOCKET NO.: 045636-5037
FULL NAME OF EIGHTH INVENTOR	Michel ARNAUD	
RESIDENCE & CITIZENSHIP	Villeurbanne, France	COUNTRY OF CITIZENSHIP FRANCE
POST OFFICE ADDRESS	63, rue Gervais Bussiére, F-69100 Villeurbanne, France	
EIGHTH INVENTOR	R'S SIGNATURE  YU'CH'D ARNAUD	DATE 2/31/20∞
FULL NAME OF ININTH INVENTOR	Michel JOLIVET	
RESIDENCE & CITIZENSHIP	Bron, France	COUNTRY OF CITIZENSHIP FRANCE
POST OFFICE ADDRESS	16, avenue des Colonnes F-69500 Bron, France	
NINTH INVENTO	R'S SIGNATURE	DATE 7/26/2003
	f Inventors Continued on attached page(s) [] Yes	X] No

1-44/(4374)3 (

Page 6 of 6